

Kinetic investigation on Effects of Cobalt ion on alkaline dissociation of Tris(1,10- Phenanthroline) iron (II) in SDS micellar media

S.Shobha Rani^{1*}, Satya guru TVSPV², V.Tejeswara Rao^{3*}

^{1*}Principal, Visakha Government Degree college for women, Visakhapatnam.

^{3*}Department of Chemistry, MVR College, Visakhapatnam, Andhra Pradesh 530045, India

²Vignana's Institute of Information Technology (A), Duvvada, Visakhapatnam-530049

ABSTRACT

Many metal ions have been found to exert electrophilic effects on the dissociation of complexes and it has been proposed that a ternary intermediate is formed involving the complex and metal ion through the severing of one of the metal ligand bonds. To investigate this aspect, the dissociation of Fe(phen)_3^{2+} and effects of metal ion like Co^{2+} in the presence of SDS micelles providing anionic micellar surface. These results confirm that the formation of such a ternary intermediate and marked micellar catalysis of the reaction.

Keywords: Spectrophotometric Determination, Co^{2+} , Tris(1,10- Phenanthroline) iron (II), Surfactant SDS.

INTRODUCTION

The study of dissociation of tris(2,2-bipyridyl)iron(II), $[\text{Fe(bipy)}_3]^{2+}$, and tris(1,10-Phenanthroline)iron(II), $[\text{Fe(Phen)}_3]^{2+}$, attracted considerable attention of workers. Basolo et al [1] carried out a detailed kinetic study of dissociation of $[\text{Fe(bipy)}_3]^{2+}$, in the aqueous medium in the presence of acid. The dissociation of the complex is accelerated by H^+ ion, the rate reaching a limiting value at $[\text{H}^+] = 1.0 \text{ mol dm}^{-3}$. Sriramam et al [2] also investigated the reaction in the aqueous acetic acid media including mineral acids and found a linear variation of rate with H^+ ion concentration reaching a limiting value at $[\text{H}^+] = 0.6 \text{ mol dm}^{-3}$ at $\mu = 0.1 \text{ mol dm}^{-3}$.

The solvent effect on the dissociation of $[\text{Fe(Phen)}_3]^{2+}$ and related complexes has been investigated in detail by Blandamer and Burgess and their co-workers [3-5]. A lot of work has been done on the kinetics of micelle-catalyzed dissociation or binding of iron(II) phenanthroline complex and its methyl substituted analogues, such as tris (3,4,7,8-tetramethyl-1,10-phenanthroline) iron(II) [6-14].



Micellar catalysis by surface-active agents or surfactants has become a subject of great significance because several aspects of this phenomenon exhibit features common with enzyme catalysis [15]. The presence of surface-active agent often brings about acceleration of reaction by 10-100 times and in a few cases 1000-10000 times [16-18].

Nucleophilic effects of anions on the complexes have been studied extensively [19-23] Subbarao et al. [24] reported that the dissociation of tris(2,2'-bipyridyl) iron(II), $[\text{Fe(bipy)}_3]^{2+}$, which is immeasurably slow in the aqueous medium in the absence of added acid, takes place with an appreciable rate reverse micellar medium and also stated that the added anions have accelerating effect on rate. Venkateswarlu and Krishna Rao investigated the Kinetics of the dissociation of tris(2,2-bipyridyl)iron(II) and tris(1,10-phenanthroline)iron(II) in the reverse micelles of Tween-85 in cyclohexane [25].

In addition to the nucleophilic effects of anions, electrophilic effects of metal ions on the complexes are also possible. Kopanica and Stara [26] reported the substitution reaction between hydrated Copper (II) and Cobalt(II) triethylenetetraminehexa acetate and the reverse

reaction, in which binuclear intermediate has been proposed. Reports are also available in the literature [27-28] that metal ions exert Lewis acid effects on complexes weakening the coordinated bond making them amenable for nucleophilic attack.

From the studies of electrophilic effects of the metal ion on the dissociation of metal complexes, Subba Rao et al [29] found kinetic evidence for the ternary intermediate involving the dissociating complex and these metal ions.



It is interesting to investigate whether these reactions are further accelerated in the micellar media in which the micelle has been oppositely charged surface to that of the metal ion and the dissociating complex and whether the reaction proceeds through mechanism involving a ternary intermediate.

To investigate this aspect, the present author has chosen the dissociation of the cationic complex $Fe(phen)_3^{2+}$ and effects of Co^{2+} metal ion in the presence of SDS micelles providing anionic micellar surface. These results are reported in this dissertation.

PREPARATION OF SOLUTIONS:

All the materials employed in this investigation were of analytical reagent grade. Solutions were prepared using water, distilled thrice in all-glass apparatus.

Tris (1,10-Phenanthroline) iron (II) :-

BDH, AnalaR sample of 1,10-Phenanthroline was used for the preparation. The Complex was prepared by mixing stoichiometric amounts (molar ratio 1:3) of ferrous ammonium sulphate (E. Merck) and 1,10-Phenanthroline in water.

Sodium dodecylsulphate (SDS):

Fluka sample of sodium dodecyl sulphate (SDS) has been used in the preparation of 0.1 mol dm^{-3}

solution. The purity of the sample was tested by determining the critical micelle concentration (cmc) ($8.0 \times 10^{-3} \text{ mol dm}^{-3}$) conductometrically.

PAN Indicator:

Dissolve 50mg of the compound in 100ml of methanol or ethanol.

Cobaltsulphate: A known volume of the cobalt solution is transferred into the conical flask add standard EDTA solution in 5 to 10ml excess followed by 10ml of acetate Buffer solution (ammonium hydroxide- ammonium chloride, p^H 10), heat to boiling and back titrate the excess of EDTA with standard copper solution using PAN as indicator.

Kinetic measurements:

The course of each of the reactions investigated has been followed by measuring the absorbance due to $[Fe(phen)_3]^{2+}$ $\lambda_{max} = 510 \text{ nm}$ ($\epsilon = 11,000 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) using Milton Roy spectronic 1201 uv-visible spectrophotometer with thermostatic arrangement.

Treatment of rate data

In all these reactions, plots of $\log A_t$ (absorbance at time 't') versus time have been found to be linear for at least 98 % of the reaction. In all the experiments, the absorbance is due to the instantaneous concentration of $[Fe(phen)_3]^{2+}$. The pseudo first order rate constants k^1 have been obtained by multiplying slope of the plot with 2.303. All the kinetic runs have been carried out in triplicate and reproducible within $\pm 5\%$.

Product Analysis

The aqueous solutions of tris (1, 10-phenanthroline) iron (II) $[Fe(phen)_3]^{2+}$ is intensely coloured and $\lambda_{max} = 510 \text{ nm}$ ($\epsilon = 11,000 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$). In the dissociation of $[Fe(phen)_3]^{2+}$ the products of dissociation are $Fe^{2+}(aq)$ and 1,10-phenanthroline. The dissociation has been established by the decrease in absorbance of the complex at $\lambda_{max} = 510 \text{ nm}$ with time, where the products of dissociation $Fe^{2+}(aq)$ and phen have no absorbance. To the product, ammonium thiocyanate was added and no red colour is developed indicating the absence of Fe(III), the products being $Fe^{2+}(aq)$ and phen, where phen = 1, 10-phenanthroline. In accord with this the UV-visible spectra of the reaction mixtures at the end

of the reaction overlap with those of 1, 10-phenanthroline at identical concentrations.

DETERMINATION OF THE CMC OF SODIUM DODECYL SULPHATE UNDER THE EXPERIMENTAL CONDITIONS EMPLOYED:

The critical micelle concentration of SDS has a value of $8.0 \times 10^{-3} \text{ mol dm}^{-3}$ in the absence of any added ions [31], but the value decreases as the ionic strength of the medium increases. The author has determined CMC at $\mu = 0.1 \text{ mol dm}^{-3}$ with respect to sodium per chlorate and in the presence of $[\text{Fe}(\text{phen})_3]^{2+} = 4.0 \times 10^{-5} \text{ mol dm}^{-3}$

by determining the surface tension of the mixture at different concentrations of SDS using stalagmometre. The value of surface tension is plotted against logarithms of SDS concentration. The decrease in surface tension with increase in [SDS] is very pronounced at SDS, the decrease in surface tension is much less pronounced. The plot has been found to yield two straight lines intersecting at $[\text{SDS}] = 1.0 \times 10^{-3} \text{ mol dm}^{-3}$ and hence, the CMC under experimental conditions has been taken to be $1.0 \times 10^{-3} \text{ mol dm}^{-3}$ (Fig 1). This result is in agreement with the CMC reported under similar experimental conditions [32-33].

Table 1: Determination of CMC

$$[\text{Fe}(\text{II})]_0 = 4.0 \times 10^{-5} \text{ mol dm}^{-3} \mu = 0.1 \text{ mol dm}^{-3} \text{Temp} = 30.0 \pm 0.1^\circ\text{C}$$

$10^{-3}[\text{SDS}]_0$ mol dm^{-3}	$10^3(\text{ Surface Tension})$ Jm^{-2}
0.0	6.48
0.4	3.97
0.6	3.71
0.8	3.49
1.0	3.29
1.2	3.27
1.5	3.24
2.0	3.19
2.5	3.11
3.0	3.11
3.5	3.11
4.0	3.11

$[\text{Fe}(\text{Phen})_3^{+2}] = 4.0 \times 10^{-5} \text{ mol dm}^{-3}$	$\text{Temp.} = 25 \pm 0.1^\circ\text{C}$
--	---




Fig. 1. Plot Graph log [SDS] Vs Surface Tension

Effect of variation of $[\text{Co}^{2+}]$ ion on rate in aqueous medium:

To investigate the effect of concentration of the Co^{+2} on the rate of dissociation of $\text{Fe}(\text{Phen})_3^{2+}$ in the aqueous medium, kinetic runs were carried out in the presence of different

concentrations of each of the metal ion, at constant ionic strength. The pseudo first order rate constants k' are presented in table 2. These results show that k' increases with increase in the metal ion concentration and shows limiting behavior at higher metal ion concentration. The plots of $1/k'$ Vs $1/[\text{metal ion}]$ are linear with positive intercepts (Fig 2).

TABLE –2: Effect of variation of $[\text{Co}^{2+}]$ on rate

$[\text{Fe}(\text{Phen})_3^{2+}] = 4.0 \times 10^{-5} \text{ mol dm}^{-3}$ Temp = $25.0 \pm 0.1^\circ\text{C}$
 $\mu = 0.01 \text{ mol dm}^{-3}$

$10^3 [\text{M}^{+2}]$ mol dm^{-3}	$10^4 \cdot k'$ Sec^{-1}
1.0	1.80
2.0	2.30
4.0	2.70
5.0	2.75
8.0	2.89
10.0	3.0

$[\text{Fe}(\text{Phen})_3^{2+}] = 4.0 \times 10^{-5} \text{ mol dm}^{-3}$ Temp = $25.0 \pm 0.1^\circ\text{C}$
 $\mu = 0.01 \text{ mol dm}^{-3}$




Fig. 2. Effect of variation of $[\text{Co}^{2+}]$ on rate

Effect of variation of metal ion concentration on rate in presence of SDS micelles:

To investigate the effect of concentration of each of the metal ion Co^{2+} on the rate of dissociation of $\text{Fe}(\text{Phen})_3^{2+}$ in the anionic micellar medium, kinetic runs were carried out in the presence of different concentrations of each of the metal ion, at constant ionic strength and at constant $[\text{SDS}]$.

The pseudo first order rate constants k' are presented in table 3. These results show that k' increases with increase in the metal ion concentration and shows limiting behavior at higher metal ion concentration. The plots of $1/k'$

Vs $1/[\text{metal ion}]$ are linear with positive intercepts (Fig 3). This shows that kinetic features of the reactions are similar in aqueous and micellar media.

$$[\text{Fe}(\text{Phen})_3^{2+}] = 4.0 \times 10^{-5} \text{ mol dm}^{-3} \quad [\text{SDS}] = 2.0 \times 10^{-3} \text{ mol dm}^{-3}$$

$$\text{Temp} = 25.0 \pm 0.1^\circ\text{C} \quad \mu = 0.01 \text{ mol dm}^{-3}$$




Fig.3. Effect of variation of $[\text{Co}^{2+}]$ on rate

Effect of Sodium dodecyl sulphate on reaction rate:

To investigate the effect of [SDS] on the rate of the reaction, kinetic runs were carried out

at constant ionic strength by varying [SDS] but keeping the concentrations of Co^{2+} constant.

The pseudo first order rate constants k' are presented in table 4.

TABLE – 4: Effect of SDS on reaction rate

$$[\text{Fe}(\text{Phen})_3^{2+}] = 4.0 \times 10^{-5} \text{ mol dm}^{-3} \quad [\text{Co}^{2+}] = 8.0 \times 10^{-4} \text{ mol dm}^{-3}$$

$$\text{Temp} = 25.0 \pm 0.1^\circ\text{C}$$

$10^3 [\text{SDS}]$ (mol dm^{-3})	$10^3 \times k$ (Sec^{-1})
3.0	3.95
4.0	4.46
5.0	4.85
6.0	5.04
8.0	5.38
10.0	5.55

The results in table 4 show that the reaction rate appreciably increases in the micellar medium. In presence of anionic SDS micelles, the rate of reaction increases by 8 and 13 times in case of Copper and Cobalt ions respectively. Further, rate constants show a tendency towards limiting behavior at higher [SDS], but no maximum is observed (Fig 4)

$$[\text{Fe(Phen)}_3^{2+}] = 4.0 \times 10^{-5} \text{ mol dm}^{-3} \quad [\text{Co}^{2+}] = 8.0 \times 10^{-4} \text{ mol dm}^{-3}$$

$$\text{Temp} = 25.0 \pm 0.1^\circ\text{C}$$




Fig. 4. Effect of SDS on reaction rate

RESULTS AND DISCUSSION

Kinetics of dissociation of $[\text{Fe(Phen)}_3]^{2+}$ in aqueous phase:

The salient kinetic features in the aqueous phase are: The reaction obeys first order kinetics with reference to Fe(Phen)_3^{2+} but fractional order kinetics with respect to the metal ion (Cu^{2+} or Co^{2+}). The plot of $1/k_{\text{obs}}$ vs $1/[\text{M}]\text{T}$ is linear with a positive intercept

The results shown that in the presence of the metal ions, the pseudo first order rate constants show tendency of limiting behavior at high

[metal ion], indicating the formation of an intermediate 1:1 add between metal ion and $[\text{Fe(phen)}_3]^{2+}$, probably one of the coordinate bonds of the complex, $[\text{Fe(phen)}_3]^{2+}$, is severed the freed nitrogen than forming a coordinate bond with the metal ion concomitantly. The metal ion in this ternary intermediate may influence further severing of the iron (II) – 1, 10-phenanthroline bonds through its electrophilic effect.

To study the stability of tris(1,10-phenanthroline) Iron(II) at pH = 6.8 (where the present investigations have been conducted) the

absorbance has been determined at various times, and the author found that there is no change in absorbance even after 6 hours due to dissociation of complex. Hence this component needs not to be considered in the analysis of rate data.

The k' versus $[M]$ plots show limiting behavior at higher metal ion concentration and the plots of $1/k_1V$ vs $1/[Metal\ ion]$ are linear with the positive slopes and intercepts.

Thus the present author has found that in addition to Ni^{2+} , Zn^{2+} , other ions like Co^{2+} , Cu^{2+} also accelerates the dissociation of $Fe(phen)_3^{2+}$ and the kinetic results prove the formation of ternary intermediate involving $Fe(phen)_3^{2+}$ - M^{2+} . The metal ion (Cu^{2+} or Co^{2+}) in the ternary intermediate exerting the electrophilic effect and accelerating the decomposition of $Fe(phen)_3^{2+}$. The authors results further established that the negative SDS micelles further accelerates the reaction and the mechanism and kinetic pattern with micellar phase are similar to those in aqueous media.

References

- [1] F Basolo, J C Hayes, H M Neumann, J. Am. Chem. Soc., 1954, 76: 3807
- [2] K Sriramam, J Sreelakshmi, L Ramadevi, and C Ramakrishna. Int. J. Chem. Kin., 1992, 24: 11.
- [3] M J Blandamer, J Burgess, S D Cope, and Digman, T. Trans. Met. Chem., 1984, 9: 347.
- [4] M J Blandamer, J Burgess, and B Clark, J. Chem. Soc., Chem. Com., 1983, 659.
- [5] J Burgess, J Chem. Soc(A), 1968, 1085.
- [6] S Tachiyashiki, and H Yamatera, Polyhedron, 1983, 2: 9.
- [7] T S Lee, I M Kolthoff, D L Leusing: J. Am. Chem. Soc., 1948, 70: 3596.
- [8] W W Brandt, D K Gullstrum: J. Am. Chem. Soc., 1952, 74: 3532.
- [9] D W Magerum: J. Am. Chem. Soc., 1957, 79: 2728.
- [10] H D Burrows, J W Ige, S A Umoh: J. Chem. Soc; Faraday Trans. I, 1982, 78: 9447.
- [11] J Ige, O O Soriyan: J. Chem. Soc. Faraday Trans. I, 1986, 82: 2011.
- [12] J Ige, J N Lambi, O O Soriyan: J. Chem. Soc, Faraday Trans. I, 1988, 1: 84.
- [13] J Ige, O Soriyan: React. Kinet. Catal Lett., 1993, 49: 419.
- [14] J Ige, O Soriyan: React. Kinet. Catal Lett., 1993, 49: 427.
- [15] I V Berezin, K Martinek and A K Yatsimirskii, Russ. Chem. Rev., 1973, 42: 787.
- [16] E H Cordes and R B Dunlop, Acct. Chem. Res., 1969, 2: 329.
- [17] E J Fendler and J H Fendler, 'Catalysis in micellar and Macro molecular systems', Academic press, New York, 1975.
- [18] A K Yatsimirskii, K Martinek and I V Berezin, Tatrahedron., 1971, 27: 2855.
- [19] S Tachiyashiki and H Yamatera, Polyhedron, 1983, 2: 9.
- [20] G Venkateswarlu, P Syamala, P V Subbarao and K Ramakrishna, J. Indian Chem. Soc., 2003, 86: 90.
- [21] P Syamala, P V Subbarao & K Ramakrishna, Indian Journal of Chemistry., 2000, 39: 643-645.
- [22] Deepa Sarkar, P V Subbarao, Gousia Begum, K Ramakrishna, Journal of Colloid and Interface Science., 2005, 288: 591-596.
- [23] Deepa Sarkar, Gousia Begum, K Ramakrishna and P V Subbarao, J. Surface Sci. Technol., 2004, 20: 9-18.
- [24] G Venkateswarlu, P Syamala, P V Subbarao & K Ramakrishna, Indian Journal of Chemistry., 2002, 41: 1410-1414
- [25] G Venkateswarlu and G S R Krishna Rao, J. Indian Chem. Soc., 2009, 86: 826-831.
- [26] M Kopanika, and V Stara, Collection Czechoslov. Chem. Comm., 1972, 37: 80.
- [27] Das, R Radha, T R Bhat, and J Sankar, J. Inorg. Nucl. Chem., 1968, 30: 1691.
- [28] M N Hughes, the Inorganic chemistry of Biological process, 2nd Edn. (Wiley), 1988, 80.
- [29] P S Phanikrishna, M. Phil Dissertation, Andhra University, 2010.
- [30] Text Book of Quantitative Chemical Analysis by I.M. Kolthoff, 4th Edition, Published by The MacMillan Company (London), p808.
- [31] C A Bunton and G Savelli, Advances in Physical organic Chemistry, 1986, 22: 216.
- [32] C A Bunton and B Wolfe, J. Am. Chem. Soc., 1973, 95: 3742.
- [33] E Pramauro, E Pelizzetti, S Diekmann and J Fahm, Inorg. Chem., 1982, 21: 2432.