

Phenotypic And Molecular Detection Of Some Virulence Genes Among Clinical Isolates Of Pseudomonas Aeruginosa And P. Fluorescence

Mustafa Sameer Midhat⁽¹⁾, Soha Maher Abd⁽²⁾, Halah abdukaliliq Awadh⁽³⁾

¹⁻³Dept.Of Biology College of Sciences – Tikrit University
Email / Mustafasamer131@gmail.com⁽¹⁾

Abstract

The study was conducted for the period from September 2021 to March 2022, in Ghazi Hariri Hospital, Yarmouk, and the Medical City. The study included collecting 115 samples from inpatients and returning patients, males, and females, of different ages. The samples included (burns, wounds, and urine) to investigate the presence of Pseudomonas aeruginosa and Pseudomonas fluorescens and study its virulence factors phenotypically and at the molecular level, as Pseudomonas aeruginosa (25) bacterial isolates were isolated from the total (115) isolates, with a percentage of (21.7%), which included (5) isolates from urine, (7) isolates from wounds, and (13) isolates from burns, which represented (20%), (28%), and (52%), respectively, from a total of 25 isolates. The results showed that 25 isolates in each of P. aeruginosa and P.fluorescence gave positive isolates for the mexB gene, as the isolates of P. aeruginosa and P.fluorescence bacteria isolated from the urine contained 25% and 20% of the MexB gene, respectively, and by 25% and 60% of wounds, while from burns it was 50% and 20%, respectively. As for the PICN gene, the study showed that the isolates of P. aeruginosa and P.fluorescence bacteria isolated from the urine contained the PICN gene by 18% and 16%, respectively, by 27% and 50% of wounds, while burns were 55% and 34%, respectively.

Also, the isolates of P. aeruginosa and P.fluorescence bacteria isolated from the urine contained the LasB gene by 22% and 25%, respectively, and by 26% and 50%, respectively, from wounds, while the percentage of burn isolates reached 52% and 25%. As for the isolates of P. aeruginosa, P.fluorescence isolated from the urine contained 20% and 19% of the AprA gene, respectively, and by 25% and 50%, respectively, from wounds, while the percentage of burn isolates reached 55% and 34%. Sequence of total bacterial isolates P. aeruginosa, P.fluorescence

Keywords: Pseudomona saeruginosa, Pseudomonas fluorescens ,mexB , PICN ,Las B. AprA gene.

Introduction

The genus Pseudomonas includes many species of bacteria everywhere, and to date, it includes 272 species isolated from many different environments, such as soil, water, air, and sediment, as well as from many

species of hosts, such as animals, plants, fungi, and algae (1). Pseudomonas aeruginosa is one of the types of Pseudomonas.spp. It is an opportunistic pathogen that possesses a group of virulence factors that make it a subject of

interest and study for researchers because of the damage it causes to human health and is responsible for some of the injuries that affect it, as it causes many acute and chronic infections such as burns, wounds, and ear infections. External, cornea, and urinary tract, and are also considered pathogens that cause acquired infections, especially in people who suffer from immunodeficiency.

Pseudomonas aeruginosa is one of the most widespread types of bacteria and its ability to cause infections among patients with wounds and burns is assisted in this by the multiple virulence factors that it possesses (1), including the formation of biofilms, the production of toxins that cause extensive damage to tissues and then enter the bloodstream and spread to the tissues of the body, as well. Because it produced virulence factors, it has high resistance to many antibiotics because it has many means of self-resistance, and it can also develop new resistance when exposed to antibacterial materials, which leads to the emergence of strains characterized by many characteristics, including multi-resistance to antibiotics. The increase in the emergence of multiple antibiotic-resistant bacteria has led many researchers to try to develop new effective antibacterial materials that have a low cost and do not cause resistance. As for the bacteria *Pseudomonas fluorescens*, it is a non-pathogenic bacterium found in soil and plants (2), and it is a chemo-organotroph for nutrition and respiration, and this bacterium is considered one of the most important factors of biological resistance (3), these bacteria are characterized by being adapted to survive in the soil or settle inside plants, and have a positive effect on both the pathosystem and growth components such as increasing plant height, wet and dry weight, and production, and are characterized by being biofilm producers (4) Enzymes produced by bacteria such as Alkaline Protease and

Elastase, where Alkaline Protease unites with Elastase to break down Collagen and Elastin. The enzyme elastase is encoded by the Las B gene, as the Las B enzyme elastase is highly efficient in analyzing proteins and penetrating the cell wall of the host (5). They have a role in analyzing many host proteins, which helps in invasion and invasion. The expression of these genes is associated with species pathogenicity, especially in skin burn patients (6). The LasB protease is encoded by the last gene. *P. aeruginosa* also has a system of efflux pumps, which are complex protein pumps that form a channel that allows materials to be pumped directly from the cytoplasm of the bacterial cell to the extracellular environment, as these pumps increase the resistance of the germ to antibiotics significantly because the effectiveness of the flow can easily exceed the flow of antibiotics. One of the largest multidrug resistance efflux pumps with a high level of expression is the MexB-OprM efflux pump, which is controlled by the regulatory genes *mix* and *mix* (7). The non-glycolytic enzyme (PLcN) Phospholipase C is encoded by the PLcN gene. *P. aeruginosa* produces two types of hemolysin: the first is called phospholipase C (PLC), which is thermally unstable and has anti-emulsifying activity, and the other is called Rhamnolipid hemolysin, which is thermally stable. Both enzymes may work together to break down lipids and lecithin, and they may also be responsible for tissue invasion via their cytotoxic effect (8).

Aim of study

1. Isolation and identification of *P. aeruginosa* and *P. fluorescens* from various clinical sources such as wounds, burns, and urinary tract infections, and studying their sensitivity to antibiotics.
2. Detection of the *lasB*, *MexB*, *AprA*, and *PLcN* genes in *P. aeruginosa* and *P. fluorescens* using polymerase chain reaction (PCR).

Materials and method

1. Collection of bacteria: 115 samples were collected from patients attending and inpatients in Ghazi Hariri, Yarmouk, and Medical City hospitals from the first of September 2021 until the end of March of the year 2020, and the collected samples included (burns, wounds, and urination) and by a sample of both sexes and of different ages, where sterile cotton swabs were used Swabs were used to take burn and wound samples. As for sputum and urine, sterile plastic containers were used. (25) isolated each of *Pseudomonas*

aeruginosa, (10) isolates from *P. fluorescens*.

2. Identification of bacteria:

Identification of bacteria: identification of the isolates by culturing on media culture including MacConkey agar, Blood agar, Cetrimide agar, *Pseudomonas* agar, and CHROMagar Orientation then identified by using biochemical tests including (oxidase and catalase test) and further identification by using an API20E system. The diagnosis was confirmed using the Vitek 2 compact system.

Table (1) Preliminary diagnostic test results for *Pseudomonas aeruginosa P. fluorescens*

<i>P. fluorescens</i> \ 10	<i>P. aeruginosa</i> 25	Properties
10	25	Fluorescin
10	25	Oxidase
0	25	Nitrate reduction
10	25	Gelatin liquefaction
10	25	Growth on cetrimide
10	0	Growth on 4C.
0	25	Growth on 42C

3. DNA isolation: A genomic DNA extraction and purification kit prepared by Promega (USA) was used to isolate genomic DNA from diagnosed bacterial isolates.

4. Detection of molecular gene: The method was used to measure the concentration and quantify the purity of DNA using a Nanodrop device, as follows: The sample to be determined

for genomic DNA was diluted 100 times using EDTA(TE) Tris dissolving solution. The optical density was measured at a wavelength of 260 nm using a UV-visible spectrophotometer and using a buffer (TE) as a control solution on the basis that a reading at a wavelength of 260 nm, which is equal to 1, is equivalent to a concentration of double-stranded DNA of 50

ng/mL. The best concentration of DNA is between (150-200) ng/ml, according to the following equation: $0.D\ 260\ \text{nm} \times \text{Dilution factor} \times 50\ \text{ug/ml} = 100$. Measurement of DNA purity:

The purity of the DNA was measured by dividing the optical density measured at the wavelength of 260 nm by the optical density measured at the wavelength of 280 nm, as the best purity of the DNA is between (1.8-2).

Results and discussion

I- Investigation of some virulence factors of *Pseudomonas* spp.

As in most types of pathogenic bacteria, *Pseudomonas* bacteria can produce many virulence factors that play an important role in the pathogenicity of these bacteria (23). The production of some virulence factors was detected in the strains of *P. fluorescens* and *P.aeruginosa*, under study, namely Biofilm Production, Haemolysin, and Lecithinase, as well as Protease production was investigated, Shows the spread of virulence factors according to the sources of isolation of these bacteria. Biofilm Production is the production of *Pseudomonas* bacteria of Biofilm mucous living membranes, which is one of the most important virulence factors possessed by bacterial strains that cause serious infections due to its lack of control due to its adhesion to many surgical tools. These results are consistent with many studies. Manandhar, (2018) mentioned that most of the 41 isolates studied by them were biofilm producers, and there was no difference in biofilm production between

the pathogenic isolates that the percentage of enzyme production was 60.5% and 69%, respectively (24).

It is clear from the current results that most of the isolates produced live mucous membranes, because the enzyme is an important virulence factor for bacteria that helps them invade, spread, damage, and resist antibiotics This current study agrees with the study by (25). Where Congo red agar and smooth surface adhesion method were used as two simple methods to examine the biofilms of *P. aeruginosa* isolates.

Molecular Detection

1- Detection of the Mex B gene

The results showed that 25 isolates in each of (*P. aeruginosa*, and *P.fluorescence*) gave positive isolates see Figure (1). The results of electrophoresis showed the presence of genetic packages with a molecular size of 151 bp compared to the size index of 100 bp in 20 isolates of *P. aeruginosa*, and 5 isolates of *P. fluorescence*, as shown in the **Figure (1)**. The results are recorded in **Table (2)**, including the distribution of the Mex B gene in the isolates under study according to the isolate sources. The results showed that the isolates of *P. aeruginosa*, *P. fluorescence* isolated from the urine contained the Mex B gene by 25% and 20%, respectively, and *P. aeruginosa* and *P.fluorescence* isolated from wounds contained the Mex B gene by 25% and 60%, respectively, while the percentage of *P. aeruginosa* and *P.fluorescence* isolated from wounds was Burns accounted for 50% and 20%, respectively, of the total bacterial isolates, *P. aeruginosa* and *P.fluorescence*.

Table (2) Distribution of Mex B gene in the isolates under study according to to isolate sources

Total	Gene distribution Mex B
-------	-------------------------

		عدد العزلات الكلي (10) P.fluorescence		P. aeruginosa عدد العزلات الكلي (25)		source of isolates
%	No.	%	No.	%	No.	
24	6	20	1	25	5	urine
32	8	60	3	25	5	Wounds
44	11	20	1	50	10	burns
100	25	100	5	100	20	• total




Figure (1): Molecular detection of Mex B gene in P. aeruginosa, P.fluorescence using PCR technology

In 2006, researchers Hearn and his group indicated that the MexB gene in *P. fluorescens* infection (MexB efflux pumps) showed importance in periplasmic domains in determining the substrate specificity of the specific substrate. The MexB flow pump has been demonstrated to discriminate and transfer substrate.

Middlemiss and Poole (2004) found several mutations within the MexB periplasmic and transmembrane domains that altered substrate selectivity, likely by affecting the tertiary structure or protein-protein interactions within the efflux complex. There is no route through which the substrate enters the central cavity, nor is there a mechanism of substrate selectivity (9). Middlemiss and Poole also identified a mutation in the outer membrane pump-membrane MexB docking protein in *P. fluorescens* that decreased the efflux activity of all antibiotics tested. This mutation in MexB was reported by Middlemiss and Poole (10).

1- Detection of the PlcN gene

The results showed that 28 isolates in each of (*P. aeruginosa*, *P. fluorescens*) gave positive isolates Figure (2). The results of electrophoresis showed the presence of genetic packages with a molecular size of 466 bp compared to the size index of 100 bp in 22 isolates of *P. aeruginosa*, and 6 isolates of *P. fluorescens*, as shown in the figure (2). The results are recorded in Table (3), which includes the distribution of the PlcN gene in the isolates under study according to the isolate sources. The results showed that the isolates of *P. aeruginosa*, *P. fluorescens* isolated from the urine contained the PlcN gene by 18% and 16%, respectively, and *P. aeruginosa* and *P. fluorescens* isolated from wounds contained the PlcN gene by 27% and 50%, respectively, while the percentage of burn isolates was 55%. And 34%, respectively, of the total bacterial isolates, *P. aeruginosa* and *P. fluorescens*.

Table (3) Distribution of PlcN gene in the isolates under study according to isolate sources

Total		Gene distribution PlcN				source of isolates
		عدد العزلات الكلي (10) P.flurosence		P. aeruginosa عدد العزلات الكلي (25)		
%	No.	%	No.	%	No.	
18	5	16	1	18	4	urine
32	9	50	3	27	6	Wounds
50	14	34	2	55	12	burns
100	28	100	6	100	22	• total

The results of the study almost agreed with the results of Endimiani and others in 2007, who found that (64%) of their isolates isolated from wounds and urine possessed this gene and it was prevalent in the isolates of urine and burns (11), and somewhat with the results of Tanya and others in 2009, who found that (72%) of Their isolates possessed this gene and it was predominant in urine (12), and the results of the current study agreed with the study of Heirmann

and others in 2012, , who recorded percentages (88%,) of their isolates isolated from different infection sites of the body (13), there are genes encoding *Pseudomonas aeruginosa* PLcN, all of which have a role in the degradation of phosphatidylcholine, and the increase in the proportions of these three genes leads to bacteremia, especially if it is present with the protease (14).






Figure (2): Molecular detection of PlcN gene in P. aeruginosa, P.fluorescens using PCR technology

2- Detection of the LasB gene

The results showed that 31 isolates in each of P. aeruginosa, and P.fluorescens) gave positive isolates see Figure (3). The results of electrophoresis showed the presence of genetic packages with a molecular size of 300 bp compared to the size index of 100 bp in 23 isolates of P. aeruginosa from a total of 25, 8 isolates of P.fluorescens from a total of 10. The results are recorded in Table (4), which includes the distribution of the last gene in the isolates under study

according to the isolate sources. The results of the study showed that the isolates of P. aeruginosa, and P.fluorescens isolated from urine, it contained the LasB gene by 22% and 25%, respectively, and P. aeruginosa and P.fluorescens isolated from wounds contained the LasB gene by 26% and 50%, respectively, while the percentage of burn isolates was 52% and 25%. %, respectively, of the total isolates of P. aeruginosa and P.fluorescens.

Table (4) Distribution of the LasB gene in the isolates under study according to isolate sources

Total		gene distribution LasB				source of isolates
		عدد العزلات الكلي (10) P.fluorescens		P. aeruginosa عدد العزلات الكلي (25)		
%	No.	%	No.	%	No.	
23	7	25	2	22	5	urine
32	10	50	4	26	6	Wounds
45	14	25	2	25	12	burns
100	31	100	8	100	23	• total

The current study agreed with that of Bradbury et al. (2010) who reported the presence of this gene in 86% of their isolates and indicated that the protease is important in burn and wound infections (15). As for the study Smith et al. (2006) that they conducted in Australia (30%) of burns and wounds have this gene (16). As

for the results of Wolska & Szweida in 2008, they recorded the presence of this gene at a rate of (100%) in their studied isolates (17), while Ghassan et al. in (2005) found the percentage of this gene in (41%) in burn (18). While the isolates of Nikbin et al. (2010), Lama et al. (2012) and Mitov et al. (2010) gave 100% of the last gene. In the

current study, we note the predominance of the gene in isolates of wounds and burns (19,20,21).

The results of Al-Shwaikh and Al-Arnawtee showed in 2019 at the University of Baghdad, where the researchers recorded

infection with *P. aeruginosa* bacteria isolated from the urine sample, where the percentage of the LasB gene was 62%, while in wounds and burns it reached 82% and 100%, respectively (5).




Figure (3): Molecular detection of LasB gene in *P. aeruginosa*, *P. fluorescens* using PCR technology

3- Detection of the AprA gene

The results showed that 26 isolates in each of (*P. aeruginosa*, and *P. fluorescens*) gave positive isolates in Figure (4). The results of electrophoresis showed the presence of genetic packages with a molecular size of 1017 bp compared with the size index of 100 bp in 20 isolates of *P. aeruginosa*, 6 isolates of *P. fluorescens*, as shown in the figure (4). The results are recorded in Table (5), which includes the distribution of the AprA gene in the isolates under study

according to the isolate sources. The results of the study showed that the isolates of *P. aeruginosa*, and *P. fluorescens* isolated from The urine contained 20% and 19% AprA gene, respectively, and *P. aeruginosa* and *P. fluorescens* isolated from wounds contained 25% and 50% AprA gene, respectively, while the percentage of burn isolates reached 55% and 34%. Sequence of total bacterial isolates *P. aeruginosa*, *P. fluorescens*.

Table (5) Distribution of the AprA gene in the isolates under study according to to isolate sources

Total		Gene distribution AprA				source of isolates
		عدد العزلات الكلي (10) P.flurosece		P. aeruginosa عدد العزلات الكلي (25)		
%	No.	%	No.	%	No.	
19	5	16	1	20	4	urine
32	8	50	3	25	5	Wounds
50	13	34	2	55	11	burns
100	26	100	6	100	20	• total

Sabharwal and authors in 2014 reported on their study, in which clinical strains were examined for the prevalence of different virulence factor genes in *P. fluorescens* and *P. aeruginosa*. The AprA gene had the lowest incidence with 16.6% of different clinical samples (22).

The large genome and genetic complexity allow *P. aeruginosa* and *P. fluorescens* to thrive in diverse environmental conditions. Multiple bacterial virulence factors influence the pathogenesis of *P. aeruginosa* infection. The combination of virulence factors expressed by each *P. aeruginosa* strain tends to determine the outcome of an infectious process. However, in clinical situations, it is often difficult to distinguish between simple colonization and infection, and there is no diagnostic tool available to assess the virulence potential of a given isolate (22).

Conclusions:

The DNA chain replication reactions were carried out using specific primers targeting the specific sequence of (Mex B, PlcN, AprA, and LasB) genes. To investigate the possession of *P. aeruginosa* bacteria, (*P. fluorescens*), of the important virulence genes that they possess.

Reference

- Govarthanan, M., Selvankumar, T., Mythili, R., Srinivasan, P., Ameen, F., AlYahya, S. A., & Kamala-Kannan, S. (2019). Biogreen remediation of chromium-contaminated soil using *Pseudomonas* sp.(RPT) and neem (*Azadirachta indica*) oil cake. *International Journal of Environmental Science and Technology*, 16(8), 4595-4600.
- Scales, B. S., Dickson, R. P., LiPuma, J. J., & Huffnagle, G. B. (2014). Microbiology, genomics, and clinical significance of the *Pseudomonas fluorescens* species complex, an unappreciated colonizer of humans. *Clinical microbiology reviews*, 27(4), 927-948.
- Kumar, B. S., & Jacob, J. (2019). Plant growth promoting rhizobacteria as a biological tool for augmenting productivity and controlling disease in agriculturally important crops-A review. *Journal of Spices & Aromatic Crops*, 28(2).
- Li, T., Wang, D., Liu, N., Ma, Y., Ding, T., Mei, Y., & Li, J. (2018). Inhibition of quorum sensing-controlled virulence factors and biofilm formation in *Pseudomonas fluorescens* by cinnamaldehyde. *International journal of food microbiology*, 269, 98-106.
- Al-Shwaikh, R. M. A., & Al-Arnawtee, A. F. (2019). Detection of *lasB* Gene of *Pseudomonasaeruginosa* Causing Different Infection. *Al-Nahrain Journal of Science*, (2), 48-52.

- 5- Islamieh, D. I., Afshar, D., & Esmaeili, D. (2019). Effect of Satureja khuzistanica essential oil (SKEO) extract on expression of lasA and lasB genes in *Pseudomonas aeruginosa*. *Iranian Journal of Microbiology*, 11(1), 55.
- 7- Pan, Y. P., Xu, Y. H., Wang, Z. X., Fang, Y. P., & Shen, J. L. (2016). Overexpression of MexAB-OprM efflux pump in carbapenem-resistant *Pseudomonas aeruginosa*. *Archives of microbiology*, 198(6), 565-571.
- 8- Kareem, S. M., Al-Kadmy, I. M., Al-Kaabi, M. H., Aziz, S. N., & Ahmad, M. (2017). *Acinetobacter baumannii* virulence is enhanced by the combined presence of virulence factors genes phospholipase C (plcN) and elastase (lasB). *Microbial pathogenesis*, 110, 568-572.
- 9- Hearn, E. M., Gray, M. R., & Foght, J. M. (2006). Mutations in the central cavity and periplasmic domain affect efflux activity of the resistance-nodulation-division pump EmhB from *Pseudomonas fluorescens* cLP6a. *Journal of bacteriology*, 188(1), 115-123.
- 10- Middlemiss, J. K., and K. Poole. 2004. Differential impact of MexB mutations on substrate selectivity of the MexAB-OprM multidrug efflux pump of *Pseudomonas aeruginosa*. *J. Bacteriol.* 186:1258–1269.
- 11- Endimani, A.;Pini,B.;Boj,A.;and Toniolo,A.(2007).Blood stream infection due top aclinical outcome associated with pathogemesis-related gene.ESCMID.European society of clinical Microbiology and Infections disease.,3(7):1164-1180.
- 12- Tanya,Z.;Strateva,K.;Guergana,B.;Petrova ,N.;and Mitov ,I. (2009) . Bulgarian Cystic fibrosis *Pseudomonas aeruginosa* isolates antimicrobial suceptibilty and nuraminidase -encoding gene distribution. .*Braz. J. Microbiol* 58(5) : 690-692.
- 13- Heirmann, G.; Yang, L.; Wu,H. ;Song, Z.; Wang,H.; Hoiby, N.; Ulrich,M.; Mollin,S.; and Doring, G. (2012).*P.aeruginosa* proteaseIV degrades surfactant protiens and inhibits surfactant host defense and biophysical functions .*J. Infection Diseases .*,202(10):1585-592.
- 14- Lama,Z.;Khairallah,M.;Sabra,A.;and Ghassan,M.(2012).Expression levels of virulence factors with up-regulated of hemolytic phospholipase C in biofilm forming *P. aeruginosa* at Tertiary care center.*J. Microbiol .*, 177(12):2266-2288.
- 15- Bradbury,L. ; Roddam, A. ;Merritt, D.; Reid,M.; and Sanre,A.(2010) .ChampionVirulence gene distribution in clinical, nosocomial and environmental isolates of *Pseudomonas aeruginosa*.*J. Medical Microbiology.*,84(5-6):499-510.
- 16- Smith,E.;Buckley,Z.;Hoffmon,L.; Argenio, D.; Miller, S.; and Moskowitz . (2006). genetic adaptation by *P. aeruginosa* to the airways of Cystic fibrosis patients .*PNAS.*,103(22).8487-84922.
- 17- Wolska,K.;and Szweda ,P .(2008).Genetic features of clinical *Ps. aeruginosa* strains. *J. Bacteriol.*, 58,3,255-260.
- 18- Ghassan,M.;Mater,M.; and Mirachaar,Z.(2005).Detection of high prevalent and potentialy virulence strain of *P. aeruginosa* from nosocominal infection in amedical center .*BMC.Microbiology.*,5,29
- 19- Nikbin,V.; Aslani, N.; and Ebrahimipour,K.(2012). Molecular identification and detection of virulence genes among *Pseudomonas aeruginosa* isolated from different infectious origins Iran *microbiol* 4(3):118-123.
- 20- Lama,Z.;Khairallah,M.;Sabra,A.;andGh assan,M.(2012).Expression levels of virulence factors with up-regulated of hemolytic phospholipase C in biofilm forming *P. aeruginosa* at Tertiary care center.*J. Microbiol .*, 177(12):2266-2288.
- 21- Mitov, I.; Strateva, T.; and Markova, B. (2010). Prevalence of virulence genes

- among bulgarian nosocomial and cystic fibrosis isolates of *Pseudomonas aeruginosa*. *Braz. J. Microbiol.* 41 : 3.
- 22- Sabharwal, N., Dhall, S., Chhibber, S., & Harjai, K. (2014). Molecular detection of virulence genes as markers in *Pseudomonas aeruginosa* isolated from urinary tract infections. *International journal of molecular epidemiology and genetics*, 5(3), 125.
- 23- Biessy, A., & Filion, M. (2021). Phloroglucinol derivatives in plant-beneficial *Pseudomonas* spp.: biosynthesis, regulation, and functions. *Metabolites*, 11(3), 182.
- 24- Manandhar, S., Singh, A., Varma, A., Pandey, S., & Shrivastava, N. (2018). Evaluation of methods to detect in vitro biofilm formation by staphylococcal clinical isolates. *BMC research notes*, 11(1), 1-6.
- 25- Lu, J., Cokcetin, N. N., Burke, C. M., Turnbull, L., Liu, M., Carter, D. A., & Harry, E. J. (2019). Honey can inhibit and eliminate biofilms produced by *Pseudomonas aeruginosa*. *Scientific reports*, 9(1), 1-13