

# EPIDEMIOLOGICAL REPORT OF SKIN LESIONS IN CHILDREN ADMITTED TO ICU SURGERY - TRAUMA

<sup>1</sup>M. Ahmadipour, <sup>2</sup>M. Ahmadijad

<sup>1</sup>Assistant Professor Pediatric Cardiologist, Faculty of Medicine Department of Pediatric Kerman University of Medical Sciences Kerman Iran.

<sup>2</sup>Associated Professor Fellowship of Critical Care Faculty of Medicine Department of Anesthesia Kerman University of Medical Sciences Kerman Iran

## Abstract

**Background:** patients who admitted in Intensive Care Units (ICUs) are susceptible to many complications. We aimed to investigate skin disorders prevalence's in critically ill surgical–trauma pediatrics during their ICU stay.

**Methods:** We recorded age, sex, Type of skin lesions, duration of ICU stay, and Pediatric index of mortality<sup>3</sup>.

We used Chi-square test ANOVA test to data analysis. Test results were evaluated at 95% confidence interval and significance level of  $P < 0.05$ .

**Results:** Our study included 632 patients (mean age of  $6.2 \pm 4$  years). 82 of the 632 (22.6%) patients were developed the skin lesions. The most common skin lesion were: Hemorrhagic lesions were (26.8%) drug eruptions (23.1%), infectious lesions (20.7%) Xeroderma, (14.6%), Pressure ulcers (9.7%), Palmoplantar keratoderma (4.8%).

The incidence of skin lesions in pediatric ICU patients was not significantly related to their age and sex ( $P\text{-Value} > 0.05$ ), the incidence of skin lesions was significantly associated with the length of ICU stay and the severity of the patient's disease. ( $P\text{-Value} > 0.05$ ).

**Conclusion:** many factors such as illness severity, duration of ICU stay, increase dermatological disorders among surgical-trauma pediatrics ICU patients.

**Keywords:** Aphthous stomatitis, Neisseria, Prevotella, Staphylococcus, Streptococcus.

## INTRODUCTION

Intensive care unit (ICU) is specialized hospital ward that provide high level treatments and monitoring's for critically ill patients such as acute respiratory distress syndrome and infectious shock (1)(2)

Acutely impairing vital organ systems and probability of life-threatening deterioration conditions requires hospitalization of the patient in the ICU(3).

Because the intensive care unit is managed in a multidisciplinary manner with the presence of specialists in different fields, the possibility of drug interactions and various drug side effects increases. (4).(5)

Also, due to underlying problems, immobility, immunodeficiency, malnutrition, etc., patients admitted to this ward are prone to various complications, including skin problems.(4)

Skin lesions are divided into primary and secondary categories, so that in the primary group are complications that cause the patient to be admitted to the ICU due to the threat to the patient's life, such as toxic epidermal necrolysis, or Versus host graft. While secondary lesions are caused by medications, immobility, or ICU infections.(7)

Some of these lesions, such as bed sores, can be prevented with proper training of the treatment staff (8) while failure to identify and treat lesions in a timely manner may be associated with increased mortality and long-term morbidity.(9)

Critically ill child's who admitted to ICU are also susceptible to various skin complication. (10) Few studies have previously been performed on the skin problems of children admitted to the ICU Therefore, in this study, we decided to provide the basis for new studies in this field by epidemiologic evaluation of skin complications in pediatrics ICU patients.

## METHODS AND MATERIALS:

In this prospective cross-sectional study, in 2020 and 2021, we evaluated all children aged 2 to 12 years who were hospitalized in the surgical-trauma intensive care unit of Kerman (southeastern Iran) in terms of the incidence of various skin lesions.

Demographic information was collected from all study participants including age, sex, diagnosis, disease severity, and length of stay in the patient's ICU and outcome. Finally, we compared the demographic findings with the prevalence of various skin lesions. To assess the severity of the disease in the first 24 hours of hospitalization, we used the online calculator of the PIM 3 pediatric index of mortality scoring system, which showed the percentage of mortality probability

Statistical analysis:

We used Chi-square test to evaluate the relationship between skin lesions and demographic findings and ANOVA test to examine the relationship between skin lesions

and findings such as length of stay in ICU, mean age of patients and severity of disease based on PIM 3 rating system. Parametric distribution values were expressed as mean standard deviation (SD). Test results were evaluated at 95% confidence interval and significance level of  $P < 0.05$ .

## RESULTS:

Out of 632 patients enrolled to the study, 361 (57%) were boys and 271 (43%) were girls. The mean age of patients was  $6.2 \pm 4$  years. The mean length of hospital stay was  $3.8 \pm 3$  days.

The most common causes of hospitalization respectively were: 34% brain trauma, 28% non-brain trauma, 15% postoperative monitoring, 10% sepsis, 7% electrolyte disturbance and 6% due to various reasons. The mean hospital stay was  $7.4 \pm 3$  days.

82 of the 632 (22.6%) patients admitted to the ICU developed skin or mucosal complications. 48 (58%) of them were boys and 34 (42%) were girls. Considering that the ratio of boys to girls hospitalized in ICU was the same, so there was no difference between the two sexes in terms of skin lesions in ICU (P-Value = 0.9)

The mean age of patients with lesions was  $7.2 \pm 4$  years, which was not significantly different from the mean age of total patients ( $6.3 \pm 4$ ) (P-Value = 0.2).

The mean length of hospital stay was  $6.2 \pm 3$  days, which was significantly longer than the mean length of total hospital stay of patients ( $3.8 \pm 3$ ) (P-Value = 0.03).

In patients with a mortality rate of more than 50% according to the pediatrics index of mortality (PMI3), skin complications were 58.5% that was significantly higher than of total patients (26.7%). (P-Value=0.001).

The most common lesions include:

- Hemorrhagic lesions were 22 (26.8%) that were created in the form of hematoma and ecchymosis following blood sampling or injection and venipuncture. In some cases,

Petechia and purpura were also seen in the dependent areas due to platelet deficiency or platelet dysfunction.

- Drug eruptions were observed in 19 patients (23.1%)
- Skin and mucosal infections in 17 patients (20.7%)
- Xeroderma, which was seen in 12 patients (14.6%) and was mostly seen on the face and limbs

Pressure ulcers were seen in 8 patients (9.7%), which respectively observed in occiput (4 cases 50%), sacrum (2 patients 25%) and elbows and heels (1 patient 12.5%).

- Palmoplantar keratoderma (4.8%)

Table1: demographic characteristics of patients

p-value	With skin lesions variable	total	
0.9	48(58%) sex	361(57%)	male
	34(42%)	271(43%)	female
0.2	7.2±4    6.2 ±4	mean age(y) ±SD	
0.03	6.2 ±3    3.8±3	Mean ICU stay(day) ±SD	
0.001	(58.5%)48	169(26.7%)	PIM 3•>50%

•Pediatric index of mortality3

Table2: frequency of different skin lesions in pediatric ICU patients

Skin lesion	No	Percent %
Hemorrhagic lesions	22	26.8
Drug eruptions	19	23.1
infections	17	20.7
xeroderma	12	14.6
Pressure ulcer	8	9.7
Palmoplantar keratoderma	4	4.8

**DISCUSSION:**

Skin lesions can be present in up to 32.9% of patients requiring ICU care, Most common skin lesions are due to critical illness or various treatments (13.2%) (11)

Our results showed that 82(22.6%) of 362 children admitted to the ICU had skin lesions. After statistical analysis, we found many factors such as: duration of ICU stay and the severity of the illness which contribute to the development of dermatological disorders in pediatric ICU patients. But there was no significant relationship between the incidence of skin lesions with demographic findings such as age and sex of children. the most common skin lesions in children admitted to ICU trauma and surgery were hemorrhagic lesions (26.8%), drug eruptions (23.1%), infections (20.7%), xeroderma (14.6%), pressure ulcer (9.7%) and palmoplantar keratoderma (4.8%). respectively.

In the study of SM Meghdadi et al, 197 out of 406 ICU patients had skin problems, the highest frequency was in the range of 21-40 years (37%) and the lowest was under 10 years (2.5%) and older than 80 years (3%). 116 of patients with skin lesions (58.9%) were male and 81 of them (41.1%) were female. (12)

Sillevis Smitt, JH et al in 2011, found that 2.3% of all ICU admitted children's (average age 30 month) had skin problems while 30% of lesions were due to ICU complications and 50% of them were related to the underlying disease.(13)

Bedia et al. reported that 9.2% of ICU patients, suffered dermatological disorders.(14) In a similar study Emre et al. reported that13.9% of the patient's need to a dermatology consultation. (15) Lee et al. found that1.2% of internal ICU patients had a dermatology consultation.(16)

Srivastava A et al study resulted 164 of 776 cases (21.13%) who admitted to ICU , had skin problems while 3.05% of them had critical lesions and 29 patients died. (17)

Prashanth G et al. after a 25-month study showed that 37.1% of children had skin complications during ICU hospitalization.(10)

In our study 22.6% of child's had skin lesions the probable reason for difference in the prevalence of skin lesions between above study is due to different Study population.

The prevalence of skin lesion in our study were: Hemorrhagic lesions were 22 (26.8%) drug eruptions (23.1%), Skin and mucosal infections (20.7%), Xeroderma, (14.6%), Pressure ulcers (9.7%),- Palmoplantar keratoderma (4.8%) while In the study of SM Meghdadi et al prevalence of dermatological problem were: hemorrhagic skin lesions (23.4%), steroid acne (22.8%) and toxic epidermal necrolysis (0.5%)

overall pressure ulcer prevalence in critically ill children is reported at 0.8% to 27%(18).Pancorbo-Hidalgo, P. L et al in an epidemiological study of pressure ulcers in pediatric ICU showed that 9.39% of patients had pressure ulcers while most of them located on the occipital region(19).

The prevalence of pressure ulcer in the study of Suddaby EC et.al was 23% in pediatric ICU patients,. Occiput ulcer was more common in critically ill (PICU) patients while diaper dermatitis was more common in the general medical-surgical population.(20)

1.2% of our patients developed pressure ulcer. it seems that strict programs for preventing pressure sores can reduce its incidence. As in the above studies, the most common site of pressure ulcer was the occiput (4 cases 50% of total pressure ulcers).

In many studies, the prevalence of drug eruptions has been estimated to be between 9.3% and 21.6%, while the most common cutaneous manifestation of drug eruption has been maculopapular rashes.(14, 21) The most prevalence of cutaneous drug reactions was reported in the study of Pektas SD et al (42.5%).(20) Which was similar to our results 45.2%. Trauma patients, need to anticonvulsants and wide spectrum, and sometimes corticosteroids, which increase the incidence of drug eruptions compared to similar studies in General PICU.

Emre et al. reported that drug reactions were more common in women, and Infectious skin lesions did not show any sex dependency(23). It may be due to estrogen effects.(22)many studied denied any association between sex and dermatological disorders (8,10).

Our study failed to show any sex-related association. We believe that the most of our patients were pre-adolescent and were deprived of the potential effects of female sex hormones.

Study of Dunnill M et al have shown that parallel to increasing ICU stay time, the incidence of dermatological problems in patients has increased. (24).

The results of George SM et al study showed that length of stay in ICU was longest for patients with acute skin failure (median 4.7 days for ICU survivors and 5.1 days for ICU non-survivors).(25)

The prevalence of skin lesions in the study of Pektas SD et al was greater in patients with ICU stay > 10 days, but no association could be found between duration of stay and type of lesion (25). We found a positive and significant relationship between length of ICU stay and prevalence of skin lesions.

We believe that with increased length of ICU stay, patients are exposed to more complications, and more drugs and various treatments, leading to a greater rate of dermatological disorders.

Some studies showed that factors such as renal failure, DM, cardiac disorders, hypothyroidism and immunosuppressive drugs were associated with an increased dermatological disorder prevalence. (26, 27)

Agrawal P et al showed Acute Physiology And Chronic Health Evaluation II (APACHE-II) score, mechanical ventilation and dialysis, are association to dermatological manifestations in critically ill patients (28)

Our results also showed a direct and significant relationship between the severity of the disease in children admitted to the ICU and the incidence of skin complications. Skin and mucosal infections was (20.7%) in our study

including wound infections, mouth candidiasis, herpes simplex and impetigo. Díaz Posada reported incidence of dermatosis was 19.59%, with a predominance of fungal diseases,(26) results of showed cutaneous cellulitis and necrotizing fasciitis were the most common manifestation of skin infection disorders in ICU.(25)

Pediatric Intensive Care Unit Patients are susceptible to nosocomial infections which may be a complication of ICU hospitalization.(29)

The development of a Surgical site infections (SSI) is dependent on multiple factors, including patient , hospital, surgeon, operation type, and surgical site characteristics, among others(27) .

Infections are the predominant etiology of skin changes in the ICU, (28)Surgical site infections (SSI) are common, occurring after up to 9 % of operations depending on surgical site(30) .Surgical site infections(SSI) range from surgical site skin wound infections to the development of sever sepsis. incidence of SSI is about 3.09%(31)

In some studies, the prevalence of wound infection in the ICU is about 18%(32)

On the other hand, patients admitted to ICU are prone to fungal infections, especially oral candidiasis, due to the use of broad-spectrum antibiotics and the intake of blood products and malnutrition.

As Haylen González Gravina et al found that 69.35% of children and adolescents with cancer had oral candidiasis.(33) the most common pathogen in many studies in ICU was *Candida* (14, 34). The high prevalence of *Candida* infections in ICU patients is due to suitable temperatures and moisture, its abundance of human flora, its ability to involve intertriginous areas, and use of broad-spectrum antibiotics and total parenteral nutrition at ICU.(35)

Our study demonstrated skin and mucosal infection prevalence of (20.7%)We believe that the high prevalence of superficial infections could be due to, disrupted skin, and surgical infectious factors in surgical patients. we also believed that mouth candidiasis is common

infection in PICU due to many predisposing factors. .

### Conclusion:

Based on the findings of the present study, skin problems are common in children admitted to the ICU. There was no relation between the prevalence of skin lesions and demographic properties (age and sex) of patients. Severity illness and length of ICU stay are the most important contributor factors to the development of dermatological disorders.

Acknowledgment: we thank the Clinical Research Development Unit, Shahid Bahonar Hospital, Kerman University of Medical Sciences, Kerman, Iran for their assistance throughout all aspects of our study.

### References

- [1] Kiblasan JIA, Alsakair IMS, Villaruz MGG, Briones GV, Garcia MDN, Fabian JP, et al. Epidemiological Study on Admissions and Discharges in the Surgical Intensive Care Unit of Misurata Central Hospital, Libya. *Journal of Health Science*. 2015;5(3):58-64.
- [2] Nesreen GE, Heba AA. Respiratory training efficacy on quality of life and functional capacity in patients with Leukemia. *J. Adv. Pharm. Educ. Res*. 2019;9(2):46-52
- [3] Kasten KR, Makley AT, Kagan RJ. Update on the critical care management of severe burns. *Journal of intensive care medicine*. 2011;26(4):223-36.
- [4] Rafiei H, Abdar ME, Amiri M, Ahmadinejad M. The study of harmful and beneficial drug interactions in intensive care, Kerman, Iran. *Journal of the Intensive Care Society*. 2013;14(2):155-8.
- [5] Al Zahrani S, Eid Alosaimi M, Alamrim AA, Alotaibi M, Almatar EA, Almanea BA. Association between knowledge and drug adherence in patients with hypertension in Saudi Arabia. *Arch Pharm Pract*. 2019;10(3):71-6.
- [6]
- [7] Delmore B, Cox J, Smith D, Chu AS, Rolnitzky L. Acute skin failure in the

- critical care patient. *Advances in skin & wound care*. 2020;33(4):192-201.
- [8] Alcântara CMP, Oliveira ELdS, Campanili TCGF, Santos RSdCS, Santos VLCdG, Nogueira PC. Prevalence and associated factors of medical adhesive-related skin injury in cardiac critical care units. *Revista da Escola de Enfermagem da USP*. 2021;55.
- [9] Dealey C, Brindle CT, Black J, Alves P, Santamaria N, Call E, et al. Challenges in pressure ulcer prevention. *International wound journal*. 2015;12(3):309-12.
- [10] Rose Ekama Ilesanmi R, Ofi BA, Adejumo PO. Nurses' knowledge of pressure ulcer prevention in Ogun State, Nigeria: results of a pilot survey. *Ostomy Wound Management*. 2012;58(2):24-32.
- [11] Prashanth G, Pai VV. A retrospective cohort study of dermatological problems observed in paediatric intensive care unit. *Journal of the European Academy of Dermatology and Venereology*. 2012;26(9):1105-8.
- [12] Awal G, Kaur T. Dermatological manifestations in the ICU: a prospective observational analysis. *Journal of the Egyptian Women's Dermatologic Society*. 2018;15(2):94-9.
- [13] Meghdadi M MM, Iraj F. *Dermatology In The Intensive Care Unit. IRANIAN JOURNAL OF DERMATOLOGY* 2000;3(11):7-13.
- [14] Smitt JHS, van Woensel JB, Bos AP. Skin lesions in children admitted to the paediatric intensive care unit: an observational study. *European journal of pediatrics*. 2011;170(10):1263-5.
- [15] Badia M, Serviá L, Casanova JM, Montserrat N, Vilanova J, Vicario E, et al. Classification of dermatological disorders in critical care patients: a prospective observational study. *Journal of critical care*. 2013;28(2):220. e1-. e8.
- [16] Kara A, Ortac E, Hapa A, Ocal S, Topeli A. Dermatological Problems and Dermatology Consultations in Intensive Care Units/Yoğun Bakım Ünitesinde Karsilasilan Dermatolojik Sorunlar ve Dermatoloji Konsultasyonu. *Dahili ve Cerrahi Bilimler Yoğun Bakım Dergisi (Journal of Medical and Surgical Intensive Care Medicine)*. 2015;6(1):1-4.
- [17] Lee KT, Lee DY. Analysis of dermatologic consultations in intensive care unit patients. *Korean Journal of Dermatology*. 2011;49(11):976-82.
- [18] Srivastava A, Mathur A, Agarwal S. Dermatological Disorders in the Intensive Care Unit: A Descriptive Study at a Tertiary Care Centre. *The Journal of the Association of Physicians of India*. 2021;69(11):11-2.
- [19] Schindler CA, Mikhailov TA, Kuhn EM, Christopher J, Conway P, Ridling D, et al. Protecting fragile skin: nursing interventions to decrease development of pressure ulcers in pediatric intensive care. *American Journal of Critical Care*. 2011;20(1):26-35.
- [20] Pancorbo-Hidalgo PL, Torra-Bou JE, Garcia-Fernandez FP, Soldevilla-Agreda JJ. Prevalence of pressure injuries and other dependence-related skin lesions among paediatric patients in hospitals in Spain. *EWMA Journal*. 2018;19(2).
- [21] Suddaby EC, Barnett S, Facticeau L. Skin breakdown in acute care pediatrics. *Pediatric nursing*. 2005;31(2):132.
- [22] Gupta H, Gupta M. Dermatological manifestations in the intensive care unit—A prospective study. *skin*. 2018;9:10.
- [23] Pektas SD, Demir AK. Prospective analysis of skin findings in surgical critically ill patients intensive care unit. *Indian journal of dermatology*. 2017;62(3):297.
- [24] Emre S, Emre C, Akoglu G, Demirseren DD, Metin A. Evaluation of dermatological consultations of patients treated in intensive care unit. *Dermatology*. 2013;226(1):75-80.
- [25] Dunnill M, Handfield-Jones S, Treacher D, McGibbon D. *Dermatology in the intensive care unit. British Journal of Dermatology*. 1995;132(2):226-35.
- [26] George SM, Harrison DA, Welch CA, Nolan KM, Friedmann PS. Dermatological conditions in intensive care: a secondary analysis of the intensive care national audit & research centre (ICNARC) case mix programme database. *Critical Care*. 2008;12(1):1-10.
- [27] Díaz Posada MC. Frecuencia y caracterización de las enfermedades dermatológicas en las unidades de cuidado intensivo de la Foscil mediante búsqueda activa y pasiva. 2019.
- [28] Pedroso-Fernandez Y, Aguirre-Jaime A, Ramos MJ, Hernández M, Cuervo M,

- Bravo A, et al. Prediction of surgical site infection after colorectal surgery. *American journal of infection control*. 2016;44(4):450-4.
- [29] Agrawal P, Peter JV, George R. Dermatological manifestations and relationship to outcomes of patients admitted to a medical intensive care unit: a study from a tertiary care hospital in India. *Postgraduate medical journal*. 2013;89(1055):501-7.
- [30] Ehlenbach WJ, Curtis JR. Noninvasive ventilation for patients near the end of life: what do we know and what do we need to know? *Critical care medicine*. 2008;36(3):1003-4.
- [31] Wilson J, Ramboer I, Suetens C, group H-Sw. Hospitals in Europe Link for Infection Control through Surveillance (HELICS). Inter-country comparison of rates of surgical site infection—opportunities and limitations. *Journal of Hospital Infection*. 2007;65:165-70.
- [32] Almogbel GT, Altokhais TI, Alhothali A, Aljasser AS, Al-Qahtani KM, Arab SF, et al. Risk Factors for Surgical Site Infections in Pediatric General Surgery: A Case–Control Study. *Journal of Pediatric Infectious Diseases*. 2021.
- [33] Elias PM, Feingold KR. Does the tail wag the dog?: role of the barrier in the pathogenesis of inflammatory dermatoses and therapeutic implications. *Archives of dermatology*. 2001;137(8):1079-81.
- [34] González Gravina H, González de Morán E, Zambrano O, Lozano Chourio M, Rodríguez de Valero S, Robertis S, et al. Oral Candidiasis in children and adolescents with cancer: Identification of *Candida* spp. *Medicina Oral, Patología Oral y Cirugía Bucal (Internet)*. 2007;12(6):419-23.
- [35] Badia M, Trujillano J, Gascó E, Casanova J, Alvarez M, León M. Skin lesions in the ICU. *Intensive care medicine*. 1999;25(11):1271-6.
- [36] Acar A, Oncül O, Kuecukardali Y, Ozyurt M, Haznedaroğlu T, Cavuşlu S. Epidemiological features of *Candida* infections detected in intensive care units and risk factors affecting mortality. *Mikrobiyoloji bulteni*. 2008;42(3):451-61